

The occurrence of *Thomomys talpoides* in the diet of the great horned owl in Garden County extends the range about 40 miles northeast of the Platte River, as noted by Imler (1945) on the basis of a single specimen caught in a snake trap at the Crescent Lake Refuge. Jones (1964) had previously dismissed the record since the specimen was not preserved. *Geomys bursarius* seems to be replacing *Thomomys* in the southwestern Panhandle due to a plowing up of upland areas (Jones, 1964), and the Crescent Lake specimens may represent a now isolated population, and one worthy of study.

REFERENCES

- Cahn, A. R., and J. W. Kemp. 1930. On the food of certain owls in east central Illinois. *Auk* 47:323-328.
- Driver, E. C. 1949. Mammal remains in owl pellets. *Amer. Midl. Nat.* 41:139-142.
- Errington, P. L. 1930. The pellet analysis method of raptor food habits study. *Condor* 32:292-296.
- Hall, E. R., and K. R. Kelson. 1959. *The Mammals of North America*. Ronald Press Co., New York, 1:xxx + 1-546+79; 2:ix + 547-1083 + 79.
- Imler, R. H. 1945. Bullsnares and their control on a Nebraska wildlife refuge. *J. Wildl. Mgmt.* 9:265-273.
- Jones, J. K. 1964. Distribution and taxonomy of mammals of Nebraska. *Univ. of Kans. Publ., Mus. Nat. Hist.*, 16:1-356.
- Korschgen, L. J., and H. B. Stuart. 1972. Twenty years of avian predator-small mammal relationships in Missouri. *J. Wildl. Mgmt.* 36:269-282.
- Moon, E. L. 1940. Notes on hawk and owl pellet formation and identification. *Trans. Kans. Acad. Sci.* 43: 457-466.



Note

CORRECTION TO CHRISTMAS BIRD COUNTS FOR NORTH DAKOTA—1971: The table appearing on pages 8 and 9, *Prairie Naturalist*, Vol. 4, No. 1, March 1972, should be corrected to delete the Gray Jay appearing under the Fargo-Moorhead Area. A number one should be added for Northern Shrike under the same Area. —Robert N. Randall, Bureau of Sport Fisheries & Wildlife, Bismarck, North Dakota 58501.