

- ? Evening Primrose, Tooth-leaved (*Oenothera serrulata*)—dry or rocky soil
 R Evening Primrose (*Oenothera nuttallii*)—sandy soil
 S Flax, Wild Yellow (*Linum rigidum*, *L. sulcatum*)
 SR Gaura, Scarlet (*Gaura coccinea*)
 S Gromwell, Wild (*Onosmodium molle*)
 R Groundsel (*Senecio* spp.)
 S Loco, Purple (*Oxytropis lambertii*)—ridges
 SR Leadplant (*Amorpha canescens*)
 SR Loosestrife, Fringed (*Lysimachia ciliata*)—low ground
 S Onion, White Wild (*Allium textile*)—stony ridges
 S Onion, Pink Wild (*Allium stellatum*)
 S Rockcress (*Arabis brachycarpa*)
 R Sage, Little (*Artemisia frigida*)—gravelly soil
 R Sage, Pasture (*Artemisia ludoviciana*)
 SR Solomon's Seal (*Smilacina stellata*)—swales
 R Spiraea, Wild (*Spiraea alba*)—swales
 SR Violet, Blue (*Viola missouriensis*)—low ground
 SR Violet, Prairie (*Viola pedatifida*)



Note

RED-BELLIED WOODPECKER IN ROLETTE COUNTY, NORTH DAKOTA. During November and December 1973 my wife DeLane and I observed a female Red-bellied Woodpecker (*Centurus carolinus*) at feeding stations on the David Disrud farm, located on the east slope of the Turtle Mountains three and a half miles northwest of Rolla. We first observed the woodpecker on November 2 and 3. Additional observations were made November 23-27 and December 14-16. Mrs. David Disrud reported that this species was first noted at the feeding stations about October 12th. The individual observed demonstrated a dominant role in interspecific encounters with other species at the feeding stations. Encounters were noted between the Red-bellied Woodpecker and Hairy and Downy Woodpeckers, Blue Jay, Black-billed Magpie, Black-capped Chickadee, White-breasted Nuthatch, and a House Sparrow.

Previous North Dakota records of Red-bellied Woodpeckers, as compiled by Robert E. Stewart, are as follows: July 1939—Family observed near Mandan by Wm. Youngworth (*Iowa Bird Life*, 14:56-57). May 1962—Jamestown (Robert E. Stewart). May 1968, 1969—Along the James River near Montpelier (Lois Haynes). The American Ornithologists' Union (1957, p. 316) lists the northern and western edges of the range of the Red-bellied Woodpecker as southeastern Minnesota, Iowa, and eastern Nebraska. Our observations thus represent the most northerly observations for this species in North Dakota.

A note of thanks is extended to Robert E. Stewart of the Northern Prairie Wildlife Research Center, Jamestown, for supplying records. —Dennis T. Disrud, *Biology Department, Minot State College, Minot 58701.*